

On the least common multiple of q -binomial coefficients

Victor J. W. Guo

Department of Mathematics, East China Normal University, Shanghai 200062, People's Republic of China

jwguo@math.ecnu.edu.cn

Abstract. In this paper, we prove the following identity

$$\text{lcm} \left(\begin{bmatrix} n \\ 0 \end{bmatrix}_q, \begin{bmatrix} n \\ 1 \end{bmatrix}_q, \dots, \begin{bmatrix} n \\ n \end{bmatrix}_q \right) = \frac{\text{lcm}([1]_q, [2]_q, \dots, [n+1]_q)}{[n+1]_q},$$

where $\begin{bmatrix} n \\ k \end{bmatrix}_q$ denotes the q -binomial coefficient and $[n]_q = \frac{1-q^n}{1-q}$. This result is a q -analogue of an identity of Farhi [Amer. Math. Monthly, November (2009)].

Keywords: least common multiple; q -binomial coefficient; cyclotomic polynomial

AMS Subject Classifications (2000): 11A07; 05A30

1 Introduction

An equivalent form of the prime number theorem states that $\log \text{lcm}(1, 2, \dots, n) \sim n$ as $n \rightarrow \infty$ (see, for example, [4]). Nair [7] gave a nice proof for the well-known estimate $\text{lcm}\{1, 2, \dots, n\} \geq 2^{n-1}$, while Hanson [3] already obtained $\text{lcm}\{1, 2, \dots, n\} \leq 3^n$. Recently, Farhi [1] established the following interesting result.

Theorem 1 (Farhi) *For any positive integer n , there holds*

$$\text{lcm} \left(\binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{n} \right) = \frac{\text{lcm}(1, 2, \dots, n+1)}{n+1}. \quad (1)$$

As an application, Farhi shows that $\text{lcm}\{1, 2, \dots, n\} \geq 2^{n-1}$ follows immediately from (1).

The purpose of this note is to give a q -analogue of (1) by using cyclotomic polynomials. Recall that a natural q -analogue of the nonnegative integer n is given by $[n]_q = \frac{1-q^n}{1-q}$. The corresponding q -factorial is $[n]_q! = \prod_{k=1}^n [k]_q$ and the q -binomial coefficient $\begin{bmatrix} M \\ N \end{bmatrix}_q$ is defined as

$$\begin{bmatrix} M \\ N \end{bmatrix}_q = \begin{cases} \frac{[M]_q!}{[N]_q! [M-N]_q!}, & \text{if } 0 \leq N \leq M, \\ 0, & \text{otherwise.} \end{cases}$$

Let lcm also denote the least common multiple of a sequence of polynomials in $\mathbb{Z}[q]$. Our main result can be stated as follows:

Theorem 2 *For any positive integer n , there holds*

$$\text{lcm} \left(\begin{bmatrix} n \\ 0 \end{bmatrix}_q, \begin{bmatrix} n \\ 1 \end{bmatrix}_q, \dots, \begin{bmatrix} n \\ n \end{bmatrix}_q \right) = \frac{\text{lcm}([1]_q, [2]_q, \dots, [n+1]_q)}{[n+1]_q}. \quad (2)$$

2 Proof of Theorem 2

Let $\Phi_n(x)$ be the n -th *cyclotomic polynomial*. The following easily proved result can be found in [5, (10)] and [2].

Lemma 3 *The q -binomial coefficient $\begin{bmatrix} n \\ k \end{bmatrix}_q$ can be factorized into*

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \prod_d \Phi_d(q),$$

where the product is over all positive integers $d \leq n$ such that $\lfloor k/d \rfloor + \lfloor (n-k)/d \rfloor < \lfloor n/d \rfloor$.

Lemma 4 *Let n and d be two positive integers with $n \geq d$. Then there exists at least one positive integer k such that*

$$\lfloor k/d \rfloor + \lfloor (n-k)/d \rfloor < \lfloor n/d \rfloor \quad (3)$$

if and only if d does not divide $n+1$.

Proof. Suppose that (3) holds for some positive integer k . Let

$$k \equiv a \pmod{d}, \quad (n-k) \equiv b \pmod{d}$$

for some $1 \leq a, b \leq d-1$. Then $n \equiv a+b \pmod{d}$ and $d \leq a+b \leq 2d-2$. Namely, $n+1 \equiv a+b+1 \not\equiv 0 \pmod{d}$. Conversely, suppose that $n+1 \equiv c \pmod{d}$ for some $1 \leq c \leq d-1$. Then $k=c$ satisfies (3). This completes the proof. ■

Proof of Theorem 2. By Lemma 3, we have

$$\text{lcm} \left(\begin{bmatrix} n \\ 0 \end{bmatrix}_q, \begin{bmatrix} n \\ 1 \end{bmatrix}_q, \dots, \begin{bmatrix} n \\ n \end{bmatrix}_q \right) = \prod_d \Phi_d(q), \quad (4)$$

where the product is over all positive integers $d \leq n$ such that for some k ($1 \leq k \leq n$) there holds $\lfloor k/d \rfloor + \lfloor (n-k)/d \rfloor < \lfloor n/d \rfloor$. On the other hand, since

$$[k]_q = \frac{q^k - 1}{q - 1} = \prod_{d|k, d>1} \Phi_d(q),$$

we have

$$\frac{\text{lcm}([1]_q, [2]_q, \dots, [n+1]_q)}{[n+1]_q} = \prod_{d \leq n, d \nmid (n+1)} \Phi_d(q). \quad (5)$$

By Lemma 4, one sees that the right-hand sides of (4) and (5) are equal. This proves the theorem. ■

3 Theorem 2 is a q -analogue of Theorem 1

In this section we will show that

$$\lim_{q \rightarrow 1} \text{lcm} \left(\begin{bmatrix} n \\ 0 \end{bmatrix}_q, \begin{bmatrix} n \\ 1 \end{bmatrix}_q, \dots, \begin{bmatrix} n \\ n \end{bmatrix}_q \right) = \text{lcm} \left(\binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{n} \right), \quad (6)$$

and

$$\lim_{q \rightarrow 1} \frac{\text{lcm}([1]_q, [2]_q, \dots, [n+1]_q)}{[n+1]_q} = \frac{\text{lcm}(1, 2, \dots, n+1)}{n+1}. \quad (7)$$

We need the following property.

Lemma 5 *For any positive integer n , there holds*

$$\Phi_n(1) = \begin{cases} p, & \text{if } n = p^r \text{ is a prime power,} \\ 1, & \text{otherwise.} \end{cases}$$

Proof. See for example [6, p. 160]. ■

In view of (4), we have

$$\lim_{q \rightarrow 1} \text{lcm} \left(\begin{bmatrix} n \\ 0 \end{bmatrix}_q, \begin{bmatrix} n \\ 1 \end{bmatrix}_q, \dots, \begin{bmatrix} n \\ n \end{bmatrix}_q \right) = \prod_d \Phi_d(1), \quad (8)$$

where the product is over all positive integers $d \leq n$ such that for some k ($1 \leq k \leq n$) there holds $\lfloor k/d \rfloor + \lfloor (n-k)/d \rfloor < \lfloor n/d \rfloor$. By Lemma 5, the right-hand side of (8) can be written as

$$\prod_{\text{primes } p \leq n} p^{\sum_{r=1}^{\infty} \max_{0 \leq k \leq n} \{ \lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor \}}. \quad (9)$$

We now claim that

$$\begin{aligned} & \sum_{r=1}^{\infty} \max_{0 \leq k \leq n} \{ \lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor \} \\ &= \max_{0 \leq k \leq n} \sum_{r=1}^{\infty} (\lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor). \end{aligned} \quad (10)$$

Let $n = \sum_{i=0}^M a_i p^i$, where $0 \leq a_0, a_1, \dots, a_M \leq p-1$ and $a_M \neq 0$. By Lemma 4, the left-hand side of (10) (denoted $\text{LHS}(10)$) is equal to the number of r 's such that $p^r \leq n$ and $p^r \nmid n+1$. It follows that

$$\text{LHS}(10) = \begin{cases} 0, & \text{if } n = p^{M+1} - 1, \\ M - \min\{i: a_i \neq p-1\}, & \text{otherwise.} \end{cases}$$

It is clear that the right-hand side of (10) (denoted $\text{RHS}(10)$) is less than or equal to $\text{LHS}(10)$. If $n = p^{M+1} - 1$, then both sides of (10) are equal to 0. Assume that $n \neq p^{M+1} - 1$ and $i_0 = \min\{i: a_i \neq p - 1\}$. Taking $k = p^M - 1$, we have

$$\lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor = \begin{cases} 0, & \text{if } r = 1, \dots, i_0, \\ 1, & \text{if } r = i_0 + 1, \dots, M, \end{cases}$$

and so

$$\sum_{r=1}^{\infty} \lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor = M - i_0.$$

Thus (10) holds. Namely, the expression (9) is equal to

$$\prod_{\text{primes } p \leq n} p^{\max_{0 \leq k \leq n} \sum_{r=1}^{\infty} (\lfloor n/p^r \rfloor - \lfloor k/p^r \rfloor - \lfloor (n-k)/p^r \rfloor)} = \text{lcm} \left(\binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{n} \right).$$

This proves (6). To prove (7), we apply (5) to get

$$\lim_{q \rightarrow 1} \frac{\text{lcm}([1]_q, [2]_q, \dots, [n+1]_q)}{[n+1]_q} = \prod_{d \leq n, d \nmid (n+1)} \Phi_d(1),$$

which, by Lemma 5, is clearly equal to

$$\frac{\text{lcm}(1, 2, \dots, n+1)}{n+1}.$$

Finally, we mention that (10) has the following interesting conclusion.

Corollary 6 *Let p be a prime number and let $k_1, k_2, \dots, k_m \leq n$, $r_1 < r_2 < \dots < r_m$ be positive integers such that*

$$\lfloor n/p^{r_i} \rfloor - \lfloor k_i/p^{r_i} \rfloor - \lfloor (n - k_i)/p^{r_i} \rfloor = 1 \quad \text{for } i = 1, 2, \dots, m.$$

Then there exists a positive integer $k \leq n$ such that

$$\lfloor n/p^{r_i} \rfloor - \lfloor k/p^{r_i} \rfloor - \lfloor (n - k)/p^{r_i} \rfloor = 1 \quad \text{for } i = 1, 2, \dots, m.$$

References

- [1] B. Farhi, An identity involving the least common multiple of binomial coefficients and its application, Amer. Math. Monthly, November (2009).
- [2] V.J.W. Guo and J. Zeng, Some arithmetic properties of the q -Euler numbers and q -Salié numbers, European J. Combin. 27 (2006), 884–895.
- [3] D. Hanson, On the product of primes, Canad. Math. Bull. 15 (1972), 33–37.
- [4] G.H. Hardy and E.M. Wright, The Theory of Numbers, 5th Ed., Oxford University Press, London, 1979.

- [5] D. Knuth and H. Wilf, The power of a prime that divides a generalized binomial coefficient, J. Reine Angew. Math. **396** (1989), 212–219.
- [6] T. Nagell, Introduction to Number Theory, Wiley, New York, 1951.
- [7] M. Nair, On Chebyshev-type inequalities for primes, Amer. Math. Monthly 89 (1982), 126–129.